arXiv:1511.09102v1 [math.CA] 29 Nov 2015

TURÁN TYPE INEQUALITIES FOR THE q-EXPONENTIAL FUNCTIONS

KHALED MEHREZ

Abstract. In this paper our aim is to deduce some sharp Turán type inequalities for the remainder $q-$exponential functions. Our results are shown to be a generalization of results which were obtained by Alzer [1].

keywords: $q-$analogue of exponential functions, Turán type inequalities.
Mathematics Subject Classification (2010) 33B10, 39B62

1. Introduction

The inequalities of the type

$$f_n(x)f_{n+2}(x) - f_{n+1}(x)^2 \geq 0$$

have many applications in pure mathematics as in other branches of science. They are named by Karlin and Szegö [6], Turán-type inequalities because the first of these type of inequalities was introduced in 1941 by P. Turán [13]. More precisely, he used some results of Szegö [12] to prove the previous inequality for $x \in (-1, 1)$, where f_n is the Legendre polynomial of degree n. Actually, the Turán type inequalities have a more extensive literature and recently the results have been applied in problems arising from many fields such as information theory, economic theory and biophysics.

Motivated by these applications, the Turán type inequalities have recently come under the spotlight once again and it has been shown that, for example, the classical Gauss and Kummer hypergeometric functions, as well the generalized hypergeometric functions, satisfy naturally some Turán type inequalities [8, 9, 10]. For deep study about this subject we refer to [2, 3, 4, 11].

This paper is organized as follows: in Section 2 we present some preliminary results and notations that will be useful in the sequel. In section 3, we investigate some Turán type inequalities for the $q-$exponential functions.

2. Notations and preliminaries

Throughout this paper, we fix $q \in (0, 1)$. We refer to [5] and [7] for the definitions, notations and properies of the q-shifted factorials and the $q-$analogue of exponential functions.

2.1. Basic symbols.

Let $a \in \mathbb{R}$, the q-Schifted factorials are defined by

$$(a;q)_0 = 1, \quad (a;q)_n = \prod_{k=0}^{n-1}(1-aq^k), \quad (a;q)_\infty = \prod_{k=0}^{\infty}(1-aq^k),$$

and we write

$$(a_1, a_2, ..., a_p; q) = (a_1; q)_n (a_1; q)_n ... (a_p; q)_n, \ n = 0, 1, 2, ...$$

Note that for $q \longrightarrow 1$ the expression $\frac{(q^a;q)_n}{(1-q)^n}$ tend to $(a)_n = a(a+1)...(a+n-1)$.

2.2. **q-analogue of exponential functions.** For $q \in (0,1)$ the $q-$analogue of exponential function are given by [5, 7]

$$e(q; z) = \sum_{n=0}^{\infty} \frac{z^n}{(q, q)_n} = \frac{1}{(z; q)_\infty}, \ |z| < 1. \tag{1}$$

and

$$E(q; z) = \sum_{n=0}^{\infty} q^{\frac{n(n-1)}{2}} \frac{z^n}{(q, q)_n} = (-z; q)_\infty = \prod_{k=0}^{\infty} \left(1 + zq^k\right), \ \ z \in \mathbb{C}. \tag{2}$$

We denote by $I_n(q; z)$ and $\mathcal{I}_n(q; z)$ the differences

$$I_n(q; z) = e(q; z) - \sum_{n=0}^{n} \frac{z^n}{(q, q)_n}, \ 0 < z < 1 \tag{3}$$

and

$$\mathcal{I}_n(q; z) = e(q; z) - \sum_{n=0}^{n} q^{\frac{n(n-1)}{2}} \frac{z^n}{(q, q)_n}, \ z > 0. \tag{4}$$

where n is a nonnegative integer.

3. Turán types inequalities for $q-$analogue of exponential functions

Theorem 1. *For every $n \in \mathbb{N}$, $q \in (0,1)$ and $0 < z < 1$, The following Turán type inequalities*

$$\frac{1 - q^{n+1}}{1 - q^{n+2}} \left(I_n(q; z)\right)^2 < I_{n-1}(q; z) I_{n+1}(q; z) < \left(I_n(q; z)\right)^2, \tag{5}$$

hlods, where $\frac{1-q^{n+1}}{1-q^{n+2}}$ is the best possible constant.

Proof. Let $n \in \mathbb{N}$ and $q \in (0,1)$ we have

$$I_{n-1}(q; z) = I_n(q; z) + \frac{z^n}{(q, q)_n} \tag{6}$$

and

$$I_{n+1}(q; z) = I_n(q; z) - \frac{z^{n+1}}{(q, q)_n}. \tag{7}$$

Thus

(8)

$$\begin{aligned}
I_{n-1}(q;z)I_{n+1}(q;z)-I_n^2(q;z) &= I_n(q;z)\left(\frac{z^n}{(q,q)_n}-\frac{z^{n+1}}{(q,q)_{n+1}}\right)-\frac{z^{2n+1}}{(q,q)_n(q,q)_{n+1}}\\
&= \frac{z^n}{(q,q)_n}\left(\frac{z^{n+1}}{(q,q)_{n+1}}+\sum_{k=n+2}^{\infty}\frac{z^k}{(q,q)_k}\right)-\frac{z^{n+1}}{(q,q)_{n+1}}I_n(z,q)-\frac{z^{2n+1}}{(q,q)_n(q,q)_{n+1}}\\
&= \sum_{k=n+2}^{\infty}\frac{z^{k+n}}{(q,q)_n(q,q)_k}-\sum_{k=n+1}^{\infty}\frac{z^{k+n+1}}{(q,q)_k(q,q)_{n+1}}\\
&= \sum_{k=n+2}^{\infty}\frac{z^{k+n}}{(q,q)_n(q,q)_k}-\sum_{k=n+2}^{\infty}\frac{z^{k+n}}{(q,q)_{k-1}(q,q)_{n+1}}\\
&= \sum_{k=n+2}^{\infty}\left[\frac{(1-q^{n+1})-(1-q^k)}{(q,q)_{n+1}(q,q)_k}\right]z^{k+n}\\
&= \sum_{k=n+2}^{\infty}\left[\frac{q^k-q^{n+1}}{(q,q)_{n+1}(q,q)_k}\right]z^{k+n}<0,
\end{aligned}$$

from which follows the right hand side inequality of (5) for $0<z<1$ and $q\in(0,1)$. Now we prove the left hand side of (5). From (6) and (7) we get the inequality

$$\left[I_n(q;z)+\frac{z^n}{(q,q)_n}\right]\left[I_n(q;z)-\frac{z^{n+1}}{(q,q)_{n+1}}\right]>\frac{1-q^{n+1}}{1-q^{n+2}}\left(I_n(q;z)\right)^2 \tag{9}$$

which is equivalent to

$$\frac{q^{n+1}-q^{n+2}}{1-q^{n+2}}\left(I_n(q;z)\right)^2>\frac{z^{2n+1}}{(q,q)_n(q,q)_{n+1}}+I_n(q;z)\left(\frac{z^{n+1}}{(q,q)_{n+1}}-\frac{z^n}{(q,q)_n}\right) \tag{10}$$

On the other hand, we get

(11)

$$\begin{aligned}
\frac{q^{n+1}-q^{n+2}}{1-q^{n+2}}\left(I_n(q;z)\right)^2 &= \left(\frac{q^{n+1}-q^{n+2}}{1-q^{n+2}}\right)\sum_{k=0}^{\infty}z^{2n+2+k}\left[\sum_{j=0}^{k}\frac{1}{(q,q)_{n+1+j}(q,q)_{n+1+k-j}}\right]\\
&> \left(\frac{q^{n+1}-q^{n+2}}{1-q^{n+2}}\right)\sum_{k=0}^{\infty}\frac{k+1}{(q,q)_{n+1}(q,q)_{n+k+1}}z^{2n+2+k}\\
&> (q^{n+1}-q^{n+2})\sum_{k=0}^{\infty}\frac{k+1}{(q,q)_{n+1}[(1-q^{n+2+k})(q,q)_{n+k+1}]}z^{2n+2+k}\\
&= (q^{n+1}-q^{n+2})\sum_{k=0}^{\infty}\frac{k+1}{(q,q)_{n+1}(q,q)_{n+k+2}}z^{2n+2+k}.
\end{aligned}$$

Using the inequality

$$\frac{q^{n+1}-q^{n+k+2}}{q^{n+1}-q^{n+2}}=\frac{1-q^{k+1}}{1-q}\leq k+1,\text{for all } k\in\mathbb{N}\text{and}q\in(0,1) \tag{12}$$

and (11) we obtain

$$\frac{q^{n+1}-q^{n+2}}{1-q^{n+2}}\left(I_n(q;z)\right)^2 > \sum_{k=0}^{\infty}\frac{\left[q^{n+1}-q^{n+k+2}\right]}{(q,q)_{n+1}(q,q)_{n+k+2}}z^{2n+2+k}. \tag{13}$$

Also, Using the inequalities

$$\frac{z^{n+1}}{(q,q)_{n+1}}I_n(q;z) = \sum_{k=0}^{\infty}\frac{z^{2n+2+k}}{(q,q)_{n+1}(q,q)_{k+n+1}} \tag{14}$$

and

$$\begin{aligned}\frac{z^n}{(q,q)_n}I_n(q;z) &= \sum_{k=0}^{\infty}\frac{z^{2n+1+k}}{(q,q)_n(q,q)_{k+n+1}}\\ &= \frac{z^{2n+1}}{(q,q)_n(q,q)_{n+1}} + \sum_{k=0}^{\infty}\frac{z^{2n+2+k}}{(q,q)_n(q,q)_{k+n+2}}\end{aligned} \tag{15}$$

we obtain

(16)

$$\begin{aligned}I_n(q;z)\left(\frac{z^{n+1}}{(q,q)_{n+1}}-\frac{z^n}{(q,q)_n}\right)+\frac{z^{2n+1}}{(q,q)_n(q,q)_{n+1}} &= \sum_{k=0}^{\infty}\left[\frac{1}{(q,q)_{n+1}(q,q)_{n+k+1}}-\frac{1}{(q,q)_n(q,q)_{n+k+2}}\right]z^{2n+2+k}\\ &= \sum_{k=0}^{\infty}\left[\frac{(1-q^{n+k+2})-(1-q^{n+1})}{(q,q)_{n+1}(q,q)_{n+k+2}}\right]z^{2n+2+k}\\ &= \sum_{k=0}^{\infty}\left[\frac{q^{n+1}-q^{n+k+2}}{(q,q)_{n+1}(q,q)_{n+k+2}}\right]z^{2n+2+k}.\end{aligned}$$

Combining (13) and (16) we get the left hand side inequality of 5. Finally, since

$$\lim_{z\longrightarrow 0}\frac{I_{n-1}(q;z)I_{n+1}(q;z)}{\left(I_n(z;q)\right)^2} = \frac{1-q^{n+1}}{1-q^{n+2}}$$

we conclude that in inequality (5) the value $\frac{1-q^{n+1}}{1-q^{n+2}}$ is the best possible constant.
So the proof of Theorem 1 is complete. ■

Theorem 2. *For all $q\in(0,1)$ and for every $n\in\mathbb{N}$ the following Turán type inequalities*

$$\left[\frac{q-q^{n+2}}{1-q^{n+2}}\right]\left(\mathcal{I}_n(q;z)\right)^2 \leq \mathcal{I}_{n-1}(q;z)\mathcal{I}_{n+1}(q;z) < \left(\mathcal{I}_n(q;z)\right)^2, \tag{17}$$

hold for all $z>0$. The value $\left[\frac{q-q^{n+2}}{1-q^{n+2}}\right]$ is the best possible constant.

Proof. Let $q\in(0,1)$, $n\in\mathbb{N}$ and $z>0$, using the inequalities

$$\mathcal{I}_{n-1}(q;z) = \mathcal{I}_n(q;z) + \frac{q^{\frac{n(n-1)}{2}}}{(q;q)_n}z^n \tag{18}$$

and

$$\mathcal{I}_{n+1}(q;z)=\mathcal{I}_n(q;z)-\frac{q^{\frac{n(n+1)}{2}}}{(q;q)_n}z^n \tag{19}$$

we obtain

(20)

$$\begin{aligned}\mathcal{I}_{n+1}(q;z)\mathcal{I}_{n-1}(q;z)-(\mathcal{I}_n(q;z))^2&=\mathcal{I}_n(q;z)\left[\frac{q^{\frac{n(n-1)}{2}}}{(q;q)_n}z^n-\frac{q^{\frac{n(n+1)}{2}}}{(q;q)_{n+1}}z^{n+1}\right]-\frac{q^{n^2}z^{2n+1}}{(q;q)_n(q;q)_{n+1}}\\&=\frac{q^{\frac{n(n-1)}{2}}}{(q;q)_n}\sum_{j=n+2}^{\infty}\frac{q^{\frac{j(j-1)}{2}}}{(q;q)_j}z^{j+n}-\frac{q^{\frac{n(n+1)}{2}}}{(q;q)_{n+1}}\sum_{j=n+1}^{\infty}\frac{q^{\frac{j(j-1)}{2}}}{(q;q)_j}z^{j+n+1}\\&=\frac{q^{\frac{n(n-1)}{2}}}{(q;q)_n}\sum_{j=n+2}^{\infty}\frac{q^{\frac{j(j-1)}{2}}}{(q;q)_j}z^{j+n}-\frac{q^{\frac{n(n+1)}{2}}}{(q;q)_{n+1}}\sum_{j=n+2}^{\infty}\frac{q^{\frac{(j-2)(j-1)}{2}}}{(q;q)_{j-1}}z^{j+n}\\&=\sum_{j=n+2}^{\infty}\frac{q^{\frac{n(n-1)+(j-1)(j-2)}{2}}\left[q^{j-1}-q^n\right]}{(q;q)_{n+1}(q;q)_j}x^{j+n}<0.\end{aligned}$$

From (20) we obtain the right side inequality of (17) for all $q\in(0,1)$ and $x>0$.

The inequality

$$\left[\frac{q-q^{n+2}}{1-q^{n+2}}\right](\mathcal{I}_n(q;z))^2\leq\mathcal{I}_{n-1}(q;z)\mathcal{I}_{n+1}(q;z) \tag{21}$$

is equivalent to

$$\left(\frac{1-q}{1-q^{n+2}}\right)(\mathcal{I}_n(q;z))^2\geq\frac{q^{n^2}z^{2n+1}}{(q;q)_n(q;q)_{n+1}}+\mathcal{I}_n(q;z)\left(\frac{q^{\frac{n(n+1)}{2}}}{(q;q)_{n+1}}z^{n+1}-\frac{q^{\frac{n(n-1)}{2}}}{(q;q)_n}z^n\right). \tag{22}$$

So

$$\frac{q^{n^2}z^{2n+1}}{(q;q)_n(q;q)_{n+1}}+\mathcal{I}_n(q;z)\left(\frac{q^{\frac{n(n+1)}{2}}}{(q;q)_{n+1}}z^{n+1}-\frac{q^{\frac{n(n-1)}{2}}}{(q;q)_n}z^n\right)= \tag{23}$$

$$= \sum_{k=n+1}^{\infty} \frac{q^{\frac{k(k-1)+n(n+1)}{2}} z^{k+n+1}}{(q;q)_{n+1}(q;q)_k} - \sum_{k=n+2}^{\infty} \frac{q^{\frac{k(k-1)+n(n-1)}{2}} z^{k+n}}{(q;q)_n(q;q)_k}$$

$$= q^{\frac{n(n+1)}{2}} \sum_{k=0}^{\infty} \frac{q^{\frac{(n+k)(n+1+k)}{2}}}{(q;q)_{n+1}(q;q)_{n+1+k}} z^{2n+2+k} - q^{\frac{n(n-1)}{2}} \sum_{k=0}^{\infty} \frac{q^{\frac{(n+k+1)(n+2+k)}{2}}}{(q;q)_n(q;q)_{n+2+k}} z^{2n+2+k}$$

$$= \sum_{k=0}^{\infty} \frac{q^{\frac{n(n-1)+(n+k)(n+k+1)}{2}}}{(q;q)_n(q;q)_{n+k+1}} \left[\frac{q^n}{1-q^{n+1}} - \frac{q^{n+k+1}}{1-q^{n+k+2}} \right] z^{2n+2+k}$$

$$= \sum_{k=0}^{\infty} \frac{q^{\frac{n(n+1)+(n+k)(n+k+1)}{2}}}{(q;q)_{n+1}(q;q)_{n+k+2}} \left(1-q^{k+1}\right) z^{2n+2+k}$$

$$= \sum_{k=0}^{\infty} \frac{q^{\frac{n(n+1)+(n+k)(n+k+1)}{2}}}{(q;q)_{n+1}(q;q)_{n+k+2}} \left(1-q^{k+1}\right) z^{2n+2+k}.$$

On the other hand, since

$$(n+j+1)(n+j) + (n+k-j+1)(n+k-j) \leq n(n+1) + (n+k)(n+k+1)$$

for all $n \in \mathbb{N}$ and $0 \leq j \leq k$, we get

(24)

$$\left(\frac{1-q}{1-q^{n+2}}\right) (\mathcal{I}_n(q;z))^2 = \left(\frac{1-q}{1-q^{n+2}}\right) \sum_{k=0}^{\infty} z^{2n+2+k} \sum_{j=0}^{k} \frac{q^{\frac{(n+j)(n+j+1)+(n+k-j)(n+k-j+1)}{2}}}{(q;q)_{n+1+j}(q;q)_{n+k-j+1}}$$

$$\geq \left(\frac{1-q}{1-q^{n+2}}\right) \sum_{k=0}^{\infty} \frac{z^{2n+2+k}}{(q;q)_{n+1}(q;q)_{n+k+1}} \sum_{j=0}^{k} q^{\frac{(n+j)(n+j+1)+(n+k-j)(n+k-j+1)}{2}}$$

$$\geq \left(\frac{1-q}{1-q^{n+2}}\right) \sum_{k=0}^{\infty} \frac{z^{2n+2+k}(k+1) q^{\frac{n(n+1)+(n+k)(n+k+1)}{2}}}{(q;q)_{n+1}(q;q)_{n+k+1}}$$

$$\geq (1-q) \sum_{k=0}^{\infty} \frac{z^{2n+2+k}(k+1) q^{\frac{n(n+1)+(n+k)(n+k+1)}{2}}}{(q;q)_{n+1}(q;q)_{n+k+2}}$$

Now, from the previous inequality , (12) and (23), we have

(25)

$$\left(\frac{1-q}{1-q^{n+2}}\right) (\mathcal{I}_n(q;z))^2 \geq \sum_{k=0}^{\infty} \frac{q^{\frac{n(n+1)+(n+k)(n+k+1)}{2}}}{(q;q)_{n+1}(q;q)_{n+k+2}} (1-q^{k+1}) z^{2n+2+k}$$

$$= \frac{q^{n^2} z^{2n+1}}{(q;q)_n(q;q)_{n+1}} + \mathcal{I}_n(q;z) \left(\frac{q^{\frac{n(n+1)}{2}}}{(q;q)_{n+1}} z^{n+1} - \frac{q^{\frac{n(n-1)}{2}}}{(q;q)_n} z^n \right).$$

So, the inequalitty (22) holds, from which follows the left side inequality of (17) for all $q \in (0,1)$ and $x > 0$.
Furthermore,

$$\lim_{z \longrightarrow 0} \frac{\mathcal{I}_{n-1}(q;z)\mathcal{I}_{n+1}(q;z)}{(\mathcal{I}_n(z;q))^2} = \frac{q-q^{n+2}}{1-q^{n+2}}.$$

The proof of Theorem 2 is completed. ■

Remark 1. *Observe also that if q tends to 1 in Theorem 2, then we get the following result: If $n \in \mathbb{N}$ and $x > 0$, then the Turán type inequality:*

$$\frac{n+1}{n+2} I_n^2(x)) < I_{n-1}(x) I_{n+1}(x)$$

is valid. We note that this inequality was proved by H. Alzer [1]

References

[1] H. Alzer, An inequality for the exponential function, Arch. Math., Vol. 55, 462-464 (1990) .

[2] H. Alzer, G. Felder, A Turán-type inequality for the gamma function, J. Math. Anal. App. 350(2009) 276182.

[3] Á. Baricz, Turán type inequalities for generalized complete elliptic integrals, Math. Z. 256(4) (2007) 895911.

[4] Á. Baricz, Turán type inequalities for hypergeometric functions, Proc. Amer. Math. Soc. 136(9) (2008) 32233229.

[5] G. Gasper, M, Rahmen:*Basic Hypergeometric Series.* Encyclopedia of Mathematics and its Application, vol. 35 (1990).

[6] S. Karlin and G. Szego, On certain determinants whose elements are orthog- onal polynomials, J. Anal. Math., 8 (1961), 1-157.

[7] T. H. Koornwinder, R. F. Swarttouw: *On q-analogues of the Hankel and Fourier transform.* Trans. Am. Math. Soc. 333, 445461 (1992).

[8] K. Mehrez, S.M. Sitnik, Proofs of some conjectures on monotonicity of ratios of Kummer and Gauss hypergeometric functions and related Turantype inequalities. arXiv:1410.6120v2 [math.CA], 2014, 8 pp.

[9] K. Mehrez, S.M. Sitnik, Inequalities for Sections of Exponential Function Series and Proofs of Some Conjectures on Monotonicity of Ratios of Kummer, Gauss and Generalized Hyper- geometric Functions. RGMIA Research Report Collection, 17(2014), Article 132, 8 pp.

[10] K. Mehrez, S.M. Sitnik, Monotonicity of Ratios of q-Kummer Con?uent Hypergeometric and q-hypergeometric Functions and associated Turán Types Inequalities. arXiv:1412.1634v1, [math.CA], 2014, 9 pp.

[11] K. Mehrez, M. ben Said, J. El Kamel, Turán types inequalities for Dunkl kernel and q-Dunkl kernel.

[12] G. Szegö, Orthogonal Polynomials, 4th ed., Colloquium Publications, vol. 23, American Mathematical Society, Rhode Island, 1975.

[13] P. Turán, On the zeros of the polynomials of Legendre, Casopis Pro Péstováni Matematiky 75 (1950), 113122.

Khaled Mehrez. Département de Mathématiques ISSATK, Kairouan 3000, Tunisia.
E-mail address: k.mehrez@yahoo.fr